

# Tips and Tricks

ACTIVE DRY YEAST AND FERMENTATION EXPLAINED TO BREWERS



THE OBVIOUS CHOICE FOR BEVERAGE FERMENTATION



**Fermentis**

LESAFFRE FOR BEVERAGES

4  
A precious tool **for you**

6  
Yeast, **what is it?**

9  
**How to rehydrate** active dry yeast?

12  
**What to be** very careful about?

15  
Yeast **characteristics**

22  
Aromas, flavors **and beer styles**

24  
**Make** your choice!

28  
Glossary

50  
Notes

# We're here to help

There are some great things happening in the world of fermented beverages. We are seeing young designers, small distilleries, craft breweries, new wine estates... There are risks; there is daring and some wonderful surprises. And as with any kind of creative endeavor, there are also disappointments. This is a virtuous model, even for the market's biggest players who are pushed to be even more inventive. This is why we want to support the efforts of those who give it a try, maybe because we share this taste for innovation and initiative. This document, we designed it for you, brewers; to offer you a tool to learn how dry yeast is produced, what essential parameters will influence your fermentations, how the Fermentis yeast strains are characterized and give useful technical tips to better manage yeast in your brewery. We sincerely hope that it will be useful to you and will help you create the beers you dream of.

*Tips and Tricks can be downloaded from our website,  
as well as other practical tips and tools.*

 **Fermentis**  
LESAFFRE FOR BEVERAGES

# A precious tool for you



Constant innovation and creativity in brewing have made the success of the craft brewing industry. Brewing a large number of beers in the same premises adds to the difficulty of yeast management, while beer quality and consistency between batches are key factors to exceed customers' expectations.


**Dry yeast is a reliable answer and the choice of numerous craft brewers around the world to achieve consistent fermentations from batch to batch.** Ready to pitch, their rehydration is a simple procedure and correct yeast counts are achieved simply by pitching a known weight of dry yeast. No propagation or in-house laboratory input is needed. The consistency of fermentations also adds the advantage of predictable fermentation output, which is essential for good planning in a busy brewery.

**Fermentis is the supplier of choice for true dried lager yeasts.** Our different strains are available from recognized sources enabling high quality lager production. A range of specialty ale yeasts has also been developed to produce ales with authentic flavor profiles and a variety of specialty beers.

Each Fermentis yeast has its own characteristics; fermentation kinetics and profile, attenuation rate, alcohol tolerance, flocculation, sedimentation, organoleptic expression...

**Better knowing our yeast range and better understanding their characteristics will allow you to get the best out of them and to adapt your brewing and fermentation conditions to brew the beer you want.**

## DRY YEAST MANUFACTURING




This diagram shows the most important steps in beer production and at which stage each ingredient enters the process. Yeast affects fermentation and subsequent steps of beer production.

**Yeast plays a key role in the release of aromas; flavors and mouth feel compounds in the finished beer.** A number of compounds will be released during fermentation and as such the yeast strain and

fermentation conditions chosen by the brewer will impact the final beer. All the elements in the brewing recipe will influence the final character and the final aromas of the beer: the water composition, the minerals, the malt bill, the choice of hops and the hopping process.

**Keep in mind that the choices made prior fermentation can also influence how the yeast reacts.**

## KEY STEPS OF BREWING





# Yeast, what is it



Yeast is a fungus capable of causing fermentation of organic and animal matter. There are several species. The best known is called *Saccharomyces cerevisiae* (from the Latin "saccharo": sugar, "myces": mushroom and "cerevisiae": "brewery"). In common parlance, we usually speaks of "brewer's yeast" or "baker's yeast".

## MICROSCOPIC PICTURE OF A YEAST CELL



Yeast, *Saccharomyces cerevisiae*, is a unicellular fungi. A *Saccharomyces cerevisiae* yeast cell measures between 5 and 50  $\mu\text{m}$ .

## Yeast, what is it?

### — LIVING CELLS

Yeast, *Saccharomyces cerevisiae*, is a unicellular fungi. They are eukaryotes with a similar cell structure to plants and animals including humans. In the presence of air, the yeasts breathe and multiply abundantly. The sugar they feed on is transformed into carbon dioxide and water. This phenomenon is accompanied by an important release of energy which allows them to grow and multiply by budding. In the absence of air, sugar is largely transformed into alcohol at the expense of the energy released.

### — SIMILAR TO A PRODUCTION FACILITY

A yeast cell could be compared to a production facility on its own. To ferment correctly it needs to be supplied with the correct raw materials to be able to produce the right compounds. When a beer recipe contains 80 to 100% of malt, its nutritional quality is sufficient for the yeast health. The yeast will metabolize sugars, amino acids and nutrients from the wort to produce **ethanol, CO<sub>2</sub>, aromas and other compounds** that will bring the final flavors to the beer.




**Don't mix-up eukaryotes and prokaryotes**

**Prokaryotes** are organisms that multiply by division and are constituted by a circular chromosome which is diffused in the cytoplasm. Example: bacteria

**Eukaryotes** are microorganisms of which genetic material is situated in a specific organ called the nucleus. Example: yeast cells, animal cells...

## YEAST METABOLISM



### — THREE SUGARS INVOLVED

Sugars are supplied by the wort. Depending on the recipe chosen for the malt bill, the quantity of sugars that can be metabolized by the yeast will vary. The three main sugars of interest for the yeast are glucose, maltose and maltotriose.

#### Glucose

Glucose is a monosaccharide; it is a single hexose and is the first sugar to be assimilated by the yeast. Glucose is a basic building block of the starch, which is a long ramified glucose chain.

#### Maltose

Maltose is a disaccharide (2 glucose units). All Fermentis brewing yeasts were selected for their high maltopermease activity. Maltopermease carries the maltose from the wort to the cytosol through the cell's membrane. Maltose is then hydrolyzed into two glucoses by intracellular maltase.

#### Maltotriose

Maltotriose is a trisaccharide sugar (3 glucose units). Not all yeasts are able to metabolize it. In theory, all bottom fermenting yeasts can assimilate maltotriose. There are some top fermenting yeasts that have this capacity too, like SafAle™ WB-06, for example.

**The result of a brew with high residual maltotriose levels will give beers with more roundness and mouthfeel, while beers with a high drinkability are those that contain no or very little residual maltotriose.**



# How to rehydrate active dry yeast



Fermentis dry yeast looks like a compact sponge composed of micro balls tightened close together. This sponge is ready to absorb the water. The yeast cells need to recover the water they lost during the drying to start fermenting. The membrane of the yeast cell after drying contains circumvolutions, after its rehydration it becomes perfectly smooth.

#### FROM DRY TO LIQUID



## How to rehydrate yeast?

### — TEMPERATURE MONITORING

- Rehydrate the dry yeast into yeast cream by sprinkling it in 10 times its own weight of sterile water or wort.
- Gently stir and leave for 30 minutes.
- Finally, pitch the resultant cream into the fermentation vessel.

The rehydration step is done in a vessel outside the fermenter. The objective is to allow the yeast to recover all its functionalities before pitching.

#### ALE YEASTS

Optimum rehydration temperature  
**25-29°C**  
(77-84°F)

#### LAGER YEASTS

Optimum rehydration temperature  
**21-25°C**  
(69-77°F)

After rehydration bacterial contamination can develop in the slurry. Follow our recommendations of maximum time between rehydration in sterile water and pitching depending on the storage temperature of the rehydrated yeast.

Stored at  
**4°C (39°F)**

pitch within  
18H


Stored at  
**20°C (68°F)**

pitch within  
6H

Stored at  
**25°C (77°F)**

pitch within  
4H

### DRY YEAST REHYDRATION



## How to rehydrate yeast?

### — WATER OR WORT?

Fermentis yeast can be rehydrated with sterile water or sterile wort, but in both cases, sterility of the rehydration environment is fundamental.

**After a first hop addition and wort boiling for at least 15 minutes, collect the volume required for rehydration and leave to cool to the required temperature. Rehydrate the yeast for 30 minutes. Pitch immediately into the tank after checking the temperature of wort, in order to avoid foam.**

## Don't forget your rehydration essentials

1

**RESPECT RECOMMENDED REHYDRATION TEMPERATURES TO ASSURE THE YEAST MEMBRANE FLUIDITY**

2

**WATER OR WORT, WHATEVER YOU CHOOSE MAKE IT STERILE**

3

**DO NOT USE CHLORINATED WATER IT WILL KILL THE YEAST**

4

**DO NOT USE DEMINERALIZED WATER**

# What to be very careful about



## What to be very careful about?

### — PITCHING RATE

Pitching at the correct level will guarantee a rapid start in fermentation. Using a low pitching rate will delay the start of the fermentation and increase the risk of contamination.

Dry yeast adds the advantage of converting a dry yeast weight to accurately know the number of viable cells pitched in the wort.

FERMENTIS YEAST DOSAGE		
ALE YEASTS	50-80 g/hl (0.06-0.10 oz/gal)	Minimum of 6E09 viable cells/g
LAGER YEASTS*	80-120 g/hl (0.10-0.16 oz/gal)	

\*Values given are for fermentation between 12-15°C (53-59°F). The yeast dosage should be increased at temperatures below 12°C (53°F), up to 200 to 300g/hl (0.26-0.40 oz/gal.) at 9°C (48°F).

### — TEMPERATURES

The recommended fermentation temperature range (refer to product packaging or specification sheets) of each strain has to be respected.

The higher the temperature is at the beginning of the fermentation, the faster the fermentation will start. Using higher temperatures for your brew will increase the ester and diacetyl formation. Also for the reduction of diacetyl it may be necessary at the end of fermentation to allow the temperature to rise.

Low temperature is required at the end of the fermentation to achieve good yeast flocculation.



**Be careful, it starts right away!**

Fermentation starts immediately, but the apparition of CO<sub>2</sub> bubbles and smell will only be perceptible after 12 to 24 hours for ale yeasts and 16 to 32 hours for lager yeasts.

### — EFFECT OF OXYGEN

Oxygen is required to assure a healthy cell multiplication. Oxygen should only be added in the first eighteen hours of fermentation. Adding oxygen later will increase aldehyde and diacetyl levels.

### — YEAST RECYCLING

Reusing yeast from a previous batch requires dedicated tanks, specific know-how and needs to be done in good hygienic conditions. Laboratory equipment and staff is required to validate the quality of the cropped yeast before pitching. As far as lager yeasts are concerned, we recommend to limit their recycling to no more than 4 to 6 times.

## Bottle and cask conditioning

Yeast is used for refermentation in bottle or in cask. If the primary objective of the method is to saturate the beer in CO<sub>2</sub>, doing a refermentation brings other benefits to the beer. First, the presence of living yeast in the bottle/cask will prevent the beer from oxidation and increase its shelf life. It will also bring mouth feel and roundness to the beer.

**When selecting a yeast for refermentation some aspects need to be considered**

- ☑ Its tolerance to higher alcohol levels
- ☑ Its aroma development capabilities
- ☑ Its sugar assimilation profiles
- ☑ Its ability to settle and stick well to the bottom of the bottle/cask

After primary fermentation, yeast is often exhausted and as such we do not recommend to use cropped yeast to make a refermentation.

**The sugar addition needs to be calculated depending on the desired carbonation of the finished beer.**

Knowing that 2g of sugar give 1g of CO<sub>2</sub> and assuming there is no CO<sub>2</sub> in the green beer, 10g of sugar per liter will need to be added to saturate the beer at 5g of CO<sub>2</sub>/l.

If the green beer already contains 2g of CO<sub>2</sub>/l, then 6g of sugar per liter have to be added.

## Yeast characteristics





Fermentis led a yeast characterization study in collaboration with a technical center\* to compare the strains between themselves in standard conditions. This study was done in EBC columns. Its purpose is to characterize each strain regarding its fermentation kinetics and attenuation, its maltotriose assimilation, its alcohol tolerance, its flocculation, its sedimentation and its aromatic profile.

\*Study conducted in collaboration with Institut Meurice - Department of Brewing Sciences and Fermentation Technology - Haute Ecole Lucia de Brouckère - Brussels, Belgium.



— FERMENTATION KINETICS AND ATTENUATION

**SafLager™ range**



SAFLAGER™ W-34/70    SAFLAGER™ S-189    SAFLAGER™ S-23


**SafAle™ range**




SAFALE™ S-04    SAFALE™ S-33    SAFALE™ T-58  
 SAFALE™ BE-256    SAFALE™ K-97    SAFALE™ US-05  
 SAFALE™ WB-06

### Other SafAle™ yeast strains


This is our selection of hyper attenuating yeasts. They present a high attenuation with low residual sugar content and allow the production of different beer styles and flavors.



— SAFALE™ BE-134 20°C/13,5°P     — SAFALE™ BE-134 25°C/13,5°P  
— SAFALE™ BE-134 30°C/13,5°P     — SAFALE™ WB-06 25°C/13,5°P



— SAFALE™ HA-18 25°C/25°P     — SAFALE™ HA-18 30°C/25°P  
— SAFALE™ HA-18 25°C/30°P     — SAFALE™ HA-18 30°C/30°P



### — MALTOTRIOSE

The table below shows the amount of remaining maltotriose in g/l after fermentation for each strain.

	MALTOTRIOSE IN G/L
SAFALE™ S-04	10
SAFALE™ K-97	2
SAFALE™ US-05	3
SAFALE™ WB-06	0
SAFALE™ S-33	12
SAFALE™ T-58	11
SAFALE™ BE-256	0
SAFALE™ BE-134	0
SAFALE™ HA-18	0
SAFLAGER™ S-23	4
SAFLAGER™ S-189	2
SAFLAGER™ W-34/70	2

### — FLOCCULATION

Flocculation is the ability of yeast cells to form aggregates. If the yeast is not remaining in the foam at the end of fermentation, a high flocculent yeast could settle down fast and give a clear beer with little cells in suspension. On the opposite, a low flocculent yeast will settle down slowly and leave the beer hazy for a longer time.

	FLOCCULATION	CLARIFICATION*	SEDIMENTATION
SAFALE™ S-04	+	-	Fast
SAFALE™ K-97	+	+	Slow
SAFALE™ US-05	+/-	+/-	Medium
SAFALE™ WB-06	-	+	Slow
SAFALE™ S-33	-	-	Medium
SAFALE™ T-58	-	-	Medium
SAFALE™ BE-256	+	-	Fast
SAFALE™ BE-134	-	-	Slow
SAFALE™ HA-18	-	-	Medium
SAFLAGER™ S-23	+	-	Fast
SAFLAGER™ S-189	+	-	Fast
SAFLAGER™ W-34/70	+	-	Fast


\*Yeast in the foam at the end of fermentation.



**Ca++**

A minimum concentration of 100 mg/l of Ca++ is required to allow good flocculation.

### — ALCOHOL TOLERANCE IN % V/V



# Aromas, flavors and beer styles

Beer yeast is able to produce or contribute to body, mouth feel, flavor and many aromas which could typically be grouped into four categories: neutral, fruity, floral and spicy. However, while the yeast strain itself will obviously play a major role; the organoleptic characteristics exhibited by a yeast strain will also largely depend on process parameters (density, fermentation temperature, pitching rate...) and beer composition. Indeed, a yeast strain will not make a beer by itself but will contribute to the elaboration of it; based on the recipe (process parameters) and together with the other raw materials; primarily water, malt and hops.

As an example; SafAle™ WB-06 expression of banana flavor will significantly depend on pitching rate and fermentation temperature.

Adjacent table will list a number of beer styles with their flavor characteristics and the recommended yeast strain(s) to achieve those.



Type of beer	Organoleptic characteristics	Suggested yeast
Weissen	Hazy, wheat base, phenolic, citrusy	WB-06
Blanche	Hazy, wheat base, refreshing, spicy	WB-06, T-58, K-97
Pils	Lager beer, blond to golden, brilliant, refreshing, drinkable, slightly crispy, medium bitterness, highly digestible, neutral, malty or gently fruity	W-34/70, S-189, S-23
Session	Blond, light body, low alcohol, hoppy, high drinkability	BE-134, K-97, US-05
Kölsch	Blond, palatable, low alcohol, low bitterness, gently fruity	K-97, US-05, S-04
IPA	Blond to amber, dry and hoppy	S-04, US-05
Triple	Blond to amber, high alcohol, malty, fruity, full body, roundness	HA-18, US-05, BE-256, S-33, K-97
Saison	Blond to amber, refreshing, very dry, low alcohol, gently acidic and yeasty, hoppy, gently saturated	BE-134, WB-06, T-58
Bitter	Blond to amber, medium body and residual sweetness balanced with high bitterness, hop character	S-33, S-04, US-05
Ales (Pale/Amber/Brown)	Blond to brown, medium alcohol content, fruity (estery), more or less malty tastes & notes, nutty, caramel	S-04, BE-256, US-05
Double	Amber - Brown/Dark, high alcohol, malty, fruity, caramel, roundness	HA-18, S-33, S-04, BE-256
Scotch	Amber to brown, full bodied, malty and lightly hopped	HA-18, S-33, S-04
Barley wine	Amber - Brown, woody, slightly saturated, maderized, stewed fruit	HA-18, S-33, T-58, BE-256, K-97
Porter	Mild to dark brown with red tint, roast malt flavor and aroma, sweet to bitter flavor, medium body, fruity esters	S-04, BE-256, US-05
Stout	Dark, creamy, smooth body, chocolate, coffee, roasted	S-33, S-04
Imperial Stout	Dark, high alcohol, hot mouthfeel, chocolate, coffee, roasted	HA-18, T-58, BE-256, US-05

Recommendations can be obtained from your distributor or from a Fermentis representative.

# Make your choice



This is our specific portfolio covering brewers needs.  
We offer you efficient and qualitative strains which will help you design the beer of your dreams.  
Let's discover their main characteristics.



Make your choice!


## — ALE OR LAGER?

Fermentis supplies 2 ranges of yeast strains.

You want to make a Lager beer? Ask for our 3 dedicated yeasts. An Ale? You can select amongst 9 strains.




\* Newcomers in our range, two new yeast strains have been partially included in this study. The **SafAle™ BE-134**, ideal for Belgian-*Saison*-style beers and the **SafAle™ HA-18**, recommended for the production of particularly high attenuating beers even for very high gravity fermentation, such as "Barley Wine".



## — DRY OR FULL-BODIED BEERS?

Find the right balance between residual sugars and final alcohol.

Almost all of our yeast strains guarantee a fairly high attenuation rate: between 80% and 90%. If you want to obtain a beer with a high attenuation and a low level of residual sugars, SafAle™ BE-256 or SafAle™ BE-134 will be the obvious choices. Likewise for high-density beers, the SafAle™ HA-18 will allow a very high attenuation. However, if you want to obtain a high level of residual sugars, SafAle™ S-33 will fit perfectly.



### CONDITIONS

This study has been set up to picture and compare the flavor and aroma characteristics of our main commercial yeast strains. All have been tested in the same standard conditions, with the lowest possible impact of other ingredients, i.e. in the most neutral conditions.

Wort: 100% 2 rows spring pils malt, 15°P / Bitterness: 25 BU with iso-alpha-acids (end of boiling) / Pitching rate: 50 g ADY/hl / Fermentation: 23°C, @Atm. P.




Make your choice!

## RESIDUAL SUGARS


Looking for yeasts which leave some specific sugars behind?

SafAle™ S-33 will leave most of the maltotriose. Conversely, SafAle™ WB-06 and SafAle™ BE-256 consume almost all of it.



## ESTERS

Some specific SafAle™ strains develop a neutral profile, while other yeasts express more fruity flavor – mainly SafAle™ BE-256 and SafAle™ WB-06.




# Baseline Flavor & Aromas\*

CHOOSE YOUR FAVORITES!

SafLager™ Yeasts

SafAle™ Yeasts



\* Sensory Analysis in Standard Conditions

\*\* Phenolic Flavors

\*\*\* Raw Material Expression Facilitated

# Glossary

A

## Alcohol By Volume (v/v)

The percentage of volume of alcohol per volume of beer. To calculate the approximate volume content apply the following method:

**Initial Gravity (°P)/2,5 = % Vol**

## Alpha-Acid Content

Measurement of the potential bitterness of hops, expressed by their percentage of alpha acids. Low: 2-6%; medium: 6-10%; high 10-14%; super > 14%.

## Attenuation

Measurement of the quantity of sugar in the wort that has been fermented by the yeast into alcohol and carbon-dioxide gas.

C

## Color

There are two different analytical methods (SRM Standard Reference Method) and EBC (European Brewery Convention) to measure the color of wort and beer. SRM units are equivalent to Lovibond degrees and are used by ASBC (American Society of Brewing Chemists).

**EBC/1,97=SRM**

D

## Density

Measurement of the weight of a solution compared with the weight of an equal volume of pure water.

## Diacetyl

Is a fermentation by-product giving "butter" off flavor. It is dismantled in the end of fermentation by the yeast. Its threshold is around 0.1 mg/L.

## Dimethyl sulphide (DMS)

An important sulphur-carrying compound originating from malt. At low levels, DMS adds a crisp character, at high levels it will add corn or cabbage flavors.

E

## Esters

Aromatic compounds generated by fermentation composed of an organic acid and an alcohol. The main esters are: **Ethyl Acetate** - aroma and fruit odor - **Isoamyl Acetate** - banana ester - and **Ethyl Hexanoate**. High fermentation yeasts are preferred for their ability to produce mixtures of particular esters.

F

## Final specific gravity

The specific gravity a beer has obtained when the fermentation is over.

I

## International Bitterness Unit (IBU)

Standard unit used to measure the concentration of bitter compounds in beer, i.e. isoalpha-acids and other related components in milligrams per liter.

M

## Malt

Barley steeped in water, germinated and dried in kilns. This process produces the enzymes necessary to convert insoluble starches to soluble substances and sugars and gives the colour to the grain transferable to beer.

## Mash - Mashing

Process of enzymatically extracting and converting malt solubles to wort, in an acid uric aqueous solution. In infusion mashing, the conversion goes through different phases: the acid rest, the protein rest, saccharification & the lauter rest.

O

## Original Gravity

Specific gravity of wort prior to fermentation. Original gravity is the measure of the total amount of dissolved solids in the wort.

P

## Plato degrees

Expresses a solution's density in grams of sucrose per 100 grams of solution. Plato degrees are measured at 20°C (68°F).

S

## Sparging

Spraying the filter cake with hot water to remove the remaining malt extract.

W

## Wort

Sweet wort is the mash extract. Bitter wort is the hopped sugar solution before pitching.



# An expert in the art of fermentation

Fermentis works with everyone in the world of beer, wine, spirits and other fermented beverages. Its range of products and services covers almost all professional requirements: from safeguarding production to expressing sensory characteristics. Business Unit of the Lesaffre Group, global key player in fermentation and yeast, Fermentis builds solutions and results upon its talented experts, visionary R&D program, industrial expertise which meets the highest international quality standards and a strong and coherent marketing and communication strategy. Its mission? Become the obvious choice for brewers, winemakers and all producers of fermented beverages, helping them express their inventiveness and creativity.



THE OBVIOUS CHOICE FOR BEVERAGE FERMENTATION



**Fermentis**

LESAFFRE FOR BEVERAGES



[www.fermentis.com](http://www.fermentis.com)

A LESAFFRE  BUSINESS UNIT